



THE CHALLENGE

You can't find colorectal cancer (CRC) early in patients who refuse to be screened. Unfortunately, 33% of average risk patients eligible for colorectal cancer screening are non-compliant with any form of CRC screening including imaging and stool-based tests.

THE COLONSENTRY SOLUTION

- ColonSentry is a 7-gene, blood-based biomarker panel that can stratify non-compliant patients according to their current, relative risk of having colorectal cancer in an average-risk population.(1-2)
- The ColonSentry test has a 95% in-office compliance rate that motivates patients to comply with CRC screening.
- Data shows that patients with elevated ColonSentry scores are likely to move forward with Colonoscopy.

THE SCIENCE BEHIND OUR TEST

- ColonSentry® uses StageZero's proprietary Sentinel Principle technology, which is based on the scientific observation that circulating blood reflects, in a detectable way, what is occurring throughout the body.
- ColonSentry uses mRNA gene expression to create a genetic signature by which a patient's risk is stratified.
- The ColonSentry test was Validated in a study of 10,000 patients in North America. (1-2)
- Negative predictive value of ColonSentry is 99.6% (1-2)
- ColonSentry has been utilized on over 100,000 patients in the United States.

WHY USE THIS TEST IN MY PRACTICE?

- Get your eligible, average-risk patients who are non-compliant screened for CRC.
- When CRC is found early, the five-year survival rate is 90% (3) Found late, the five-year survival rate drops to 14%.(3)
- Young adults with colon cancer have a statistically higher risk of advanced disease at the time of operation.(7)

PATIENT SELECTION CRITERIA

- For use on men and women age 45 and over who have yet to comply with Colorectal Cancer Screening.

Contact Us Today to Get Started

1-855-420-7150

clientrelations@stagezerols.com

TEST INTERPRETATION

- If the ColonSentry score is elevated, there is an increased probability of the patient having colorectal cancer.(1-2)
- Patients with elevated ColonSentry scores should be encouraged to have a colonoscopy.
- Patients with average ColonSentry scores should continue to comply with USPSTF guidelines.

Contact Us Today to Learn More


1-855-420-7150

clientrelations@stagezerols.com



ColonSentry®
LABORATORY RESULTS

Patient	Name:	Phone #:	Patient ID #:	Specimen	Collection Time:	Specimen ID:	Provider	Requesting Provider:	
	Fasting Status:	Gender:	Birthdate:		Age:	Collection Date:		Report Type:	Client ID:
	Height:	Weight:	BMi:		Prev. BMi:	Received Date:		Report Date:	



Test Description

Test results are reported with a 95% CI (Confidence Interval). The ColonSentry® test measures the expression of 7 gene biomarkers in whole blood to help determine a patient's Current Risk for having colorectal cancer (CRC) relative to the current risk in an average risk population.

Gene Expression

Gene	Cycle Threshold
ANXA3* (ΔCt)	1.9
CLEC4D* (ΔCt)	2.5
TNFAIP6* (ΔCt)	1.8
LMNB1* (ΔCt)	1.8
PRRG4* (ΔCt)	0.9
VNN1* (ΔCt)	1.5
IL2RB (Ct)	24.5

* Difference with respect to IL2RB

Clinical Recommendations

If the results indicate that the patient's current risk is elevated (2X or more), the patient should be referred for further evaluation with procedures such as colonoscopy. Average risk scores for the ColonSentry® test do not rule out colon cancer. When caught early, colon cancer is both treatable and beatable.

Average Risk: recommendation** for FIT, FIT-DNA, flexible sigmoidoscopy, CT colonography, or colonoscopy.
Elevated Risk: recommendation** for colonoscopy.

** Screening for Colorectal Cancer US Preventive Services Task Force Recommendation Statement. JAMA Jun 21, 2016; 315(23):2564-2575

What to do next?

According to USPSTF*, an average risk patient is a man or woman who is at least 50 years old, is asymptomatic for CRC, has no personal history of benign colorectal polyps, colorectal adenomas, CRC or inflammatory bowel disease, and does not have a first degree relative with CRC. The presence of conditions such as pancreatic cancer, systemic sclerosis or CLL might affect the risk score.

Screening for colorectal cancer reduces mortality through detection and treatment of early-stage cancer and detection and removal of adenomatous polyps. The degree to which each of these mechanisms contributes to a reduction in mortality is unknown, although it is likely that the largest reduction in colorectal cancer mortality during the 10 years after initial screening comes from the detection and removal of early-stage cancer. Colonoscopy is a necessary step in any screening program that reduces mortality from colorectal cancer.***

*** Screening for Colorectal Cancer US Preventive Services Task Force Recommendation Statement. JAMA Jun 21, 2016; 315(23):2564-2575

Disclaimer

This test is not recommended for patients that have previous history of colorectal cancer, or pre-cancerous (e.g. adenomatous) polyps, or familial or inherited colon polyp syndromes, or inflammatory bowel disease, or have received chemotherapy and/or radiation.

1. Liew CC, Ma J, Tang HC, et al. The peripheral blood transcriptome dynamically reflects system wide biology: a potential diagnostic tool. J Lab Clin Med. 2006 Mar;147(3):126-32.
2. Marshall KW, Mohr S, Khattabi FE, et al. A blood-based biomarker panel for stratifying current risk for colorectal cancer. Int J Cancer. 2010 Mar 1;126(5):1177-86.
3. Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Cancer Stat Facts: Colorectal Cancer, National Cancer Institute, DCCPS, Surveillance Research Program, released December 2018. Underlying mortality data provided by NCHS (www.cdc.gov/nchs).
4. Chao S, Ying J, Liew G, et al. Blood RNA biomarker panel detects both left- and right-sided colorectal neoplasms: a case- control study. J Exp Clin Cancer Res. 2013 Jul 23;32:44.
5. Yip KT, Das PK, Suria D, et al. A case- controlled validation study of a blood- based seven-gene biomarker panel for colorectal cancer in Malaysia. J Exp Clin Cancer Res. 2010 Sep 16;29:128.
6. Han M, Liew CT, Zhang HW, et al. Novel blood-based, five-gene biomarker set for the detection of colorectal cancer. Clin Cancer Res. 2008 Jan 15;14(2):455-60.
7. Moghadamyeghaneh Z, Alizadeh RF, Phelan M, et al. Trends in Colorectal Cancer Admissions and Stage at Presentation-Impact of Screening. Surgical endoscopy. 2016;30(8):3604-3610. doi:10.1007/s00464-015-4662-3.
8. Chao S, Pilcz T, Stamatou, D, et al. Stability of The ColonSentry® Colon Cancer Risk Stratification Test. Int J of Dis Markers. 2019 DOI: 10.29011/IJDM -101. 100001